

# Transitional to Final Dentures: A Detailed Process for the Fabrication of Complete Dentures—Part 1



Lorin F. Berland, DDS  
Private Practice  
Dallas, Texas  
Phone: 214.999.0110  
Email: drberland@dallasdentalspa.com  
Web site: www.dallasdentalspa.com

**T**hanks to recent exposure in the media, advances in technology, and professional enthusiasm, there is a new public awareness of esthetic procedures. Television shows such as *Extreme Makeover* and *The Swan* focus mainly on plastic surgery; when esthetic dentistry is addressed, the emphasis is on whitening and porcelain veneers. Unfortunately, this kind of phenomenon has virtually ignored the denture-wearing public. Advancements in denture technology have made dentures better than ever. Dentists who provide patients with quality dentures can make dramatic improvements in esthetics and function, as well as overall improvement of their lives.

Approximately 25% of Americans at age 65 are edentulous and, at 75, approximately 44% are edentulous. About 17% of the adult population (30 million people) are edentulous in at least one arch.<sup>1</sup> On average, most of the author's new denture patients have worn the same dentures for far longer than the recommended 3 to 5 years. These dentures are typically more than 10 years old.

A denture that does not fit or teeth that do not function effectively can make it difficult to chew food properly. As a result, many nutritious and harder-to-chew foods are missing from the diet. In these

## Abstract

Complete denture fabrication and esthetics has been essentially overlooked with the arrival of cosmetic dentistry and dental implants. The procedures in diagnosis and analysis for complete immediate denture rehabilitation and the techniques involved are discussed in this article. Before the creation of maxillary prosthesis, a complete extra- and intra-oral examination should be performed. Along with impressions of the existing dentition and soft tissue, diagnostic occlusal bite registration is performed before processing the final prostheses. This article is just Part 1 in a two-part series. In this article, treatment of a patient with periodontal disease, failing dentition, and an old, ill-fitting mandibular denture was restored with an immediate complete maxillary denture. Part 2 follows next month with the restoration of the oral complex with new complete maxillary and mandibular dentures. Photographs illustrate the removal of the remaining dentition, placing a soft denture liner, and creating an esthetic smile.

## Learning Objectives

After reading this article, the reader should be able to:

- explain failing dentition and the treatment plan for complete immediate dentures.
- make use of proper diagnostic methods with evidence-based dentistry before treatment to assist in the fabrication of an immediate denture prosthesis.
- use and place current dental materials properly for pre- and post-insertion of the denture prosthesis.

cases, digestion and general health can be negatively affected. Furthermore, old, ill-fitting dentures can even create the illusion of an older look. Obviously this can have an adverse effect on a person's emotional well-being, and their self-confidence.

Oral structures change naturally over time. Even a denture that started out fitting well will not fit the same way it used to. One set of dentures is not a permanent solution. Dentures eventually need to be relined, rebased, or replaced, just like everything else in life. Nothing

lasts forever, and neither do dentures. Dentures that no longer fit properly can cause friction in the soft tissues, which could lead to sores, which in turn are prone to infections. An older denture is also more likely to harbor more bacteria. Recent studies have shown that disinfectants will not eliminate microorganisms such as *Escherichia coli*, *Staphylococcus aureus*, and *Candida albicans*.<sup>2</sup> The best way to eliminate the microorganisms from an old denture is to make a new one.<sup>3</sup>

## Oral Cancer

According to the American Cancer Society, oral cancer is the sixth most common cancer and accounts for almost 3% of all cancers diagnosed. There are an estimated 30,000 new cases reported annually in the United States alone.<sup>4</sup> According to the Oral Cancer Foundation, 8,000 Americans die of oral cancer each year.<sup>5</sup> The vast majority of oral cancer occurs in people older than 45, with men twice as likely as women to develop the disease.<sup>6</sup> The use of tobacco and alcohol are important risk factors for oral epithelial dysplasia. This is a histopathological diagnosis that is associated with an increased risk of oral cancer; and denture patients may be the most at risk because old dentures that no longer fit irritate the tissue.



Figure 1—Preoperative image.



Figure 2—Preoperative smile.



Figure 3—Preoperative maxillary occlusal.



Figure 4—Preoperative retracted view.



**Figure 5**—Preoperative retracted view right lateral.



**Figure 6**—Preoperative retracted view left lateral.



**Figure 7**—Mandibular edentulous occlusal view.



**Figure 8**—Cameo surface of existing denture.



**Figure 9**—Intaglio surface of existing denture with Sea-Bond.

Therefore, the Oral Cancer Foundation recommends that all denture patients get regular oral cancer screenings.<sup>7</sup>

### A Clinical Case

This article is the first of two parts. Part 1 details the transition from the partially edentulous maxilla to an immediate transitional denture. Part 2 will appear next month with the restoration of the oral complex with new complete maxillary and mandibular dentures.

### The Patient

For many, the Internet has become the primary source for accurate, unbiased information. This energetic 53-year-old man found the author's practice through [www.denturewearers.com](http://www.denturewearers.com), specifically the "Find Your Local Denture Doctor" section. He was concerned about his "very loose teeth" and he wanted a better-looking smile. He had not been to a dentist in more than 10 years when his lower teeth were extracted and replaced with a lower denture.

The extraoral examination showed no locking, deviation, restriction of movement, crepitus of the temporomandibular joint, or pain on palpation<sup>1</sup> (Figure 1). He had a protrusive profile with an Angle class III malocclusion. His facial form was ovoid and brachycephalic.<sup>2</sup> His musculature and functional tone were within normal limits. He had a support-

# Tescera<sup>TM</sup> ATL<sup>TM</sup>


## Indirect Composite Restorative System

# Pearls of Wisdom




“Conservative dentistry at it's best!”

**- Dr. Lou Graham**  
Private Practice,  
University of Dental Professionals,  
Co-Founder Dental Team Concepts



“I have been using the Tescera ATL for over two years with outstanding results. When restoring inlays and onlays and for fiber reinforced bridges the superior aesthetics, high strength, excellent wear and abrasion resistance make it my first choice for conservative, bonded laboratory fabricated restorations.”

**- Dr. Howard Strassler**  
University of Maryland



“The Tescera ATL system has renewed my confidence in placing resin onlays. The system puts strength, aesthetics and conservation of tooth structure all in a single package.”




**- Dr. Robert E. Rada**  
Private Practice, La Grange, IL  
Clinical Assistant Professor, at the  
University of Illinois at Chicago (UIC)

Circle 22 on Reader Service Card

**For Information on Tescera or Training**  
please contact Bisco at  
**800.247.3368 • [tescera@bisco.com](mailto:tescera@bisco.com) • [www.bisco.com](http://www.bisco.com)**



Bringing Science to the Art of Dentistry MC-1088LTC



Figures 10A through 10E—Preoperative periapical radiographs.



Figure 11—Digital image of proposed treatment.



Figure 12—Calipers in place to assist in placement of canines.




Figure 13—Papillometer in place to assist in vestibular depth.



Figure 14—Hydro-Cast before placement and trimming.

ed lip and low smile line (Figure 2).

Intraorally, his soft tissues were inflamed and irritated. Fortunately, he did not have any signs of oral carcinoma. Most notably, he had advanced and generalized periodontitis in the maxillary arch (Figure 3). These loose and unattractive teeth were his chief concern. He wore an old, ill-fitting mandibular denture (Figures 4 through 6). He was missing teeth Nos. 1 through 3 and 14 through 16. Tooth No. 8 was restored with a 7/8 gold crown. He was edentulous in the mandibular arch. This is classified as a favorable class I arch (Figure 7). His mandibular denture was past its prime, loose, and discolored. In fact, he always wore it with a Sea-Bond insert (Combe Incorporated) (Figures 8 and 9).

Radiographically, he had normal bone trabeculation in the mandibular arch, but suffered from advanced generalized periodontitis in the maxilla. This resulted in excessive bone loss (Figures 10A through 10E).

### Treatment Options

The challenges associated with treating this patient included generalized periodontitis in the upper arch with severe horizontal bone loss around the remaining dentition, edentulous areas, a skeletal class III malocclusion, and, most importantly, the patient's expecta-

tions. The treatment goals were removing the remaining infected dentition, providing for adequate function and phonetics, restoring an ideal occlusal plane to prevent more degenerative changes, and creating an esthetic smile. The treatment options were extractions and immediate transitional denture followed by conventional dentures, implant-supported maxillary and mandibular prostheses, a combination of both, or nothing. After dis-

**T**he treatment options were extractions and immediate transitional denture followed by conventional dentures, implant-supported maxillary and mandibular prostheses, a combination of both, or nothing.

cussing the options with the patient, the treatment plan was formulated, beginning with the fabrication of an immediate transitional maxillary denture. After an appropriate amount of healing time and adjustments, treatment will continue with new upper and lower dentures (Part II).

When fabricating an immediate denture, teeth and gingival shades, as well as lengths, should be selected based on the individual's characteristics and desires. Digital photographs can be most helpful at this

stage in patient communication. Esthetic imaging can also be a useful tool in helping patients understand needed dental treatment. This kind of visualization help patients "see" what esthetic dentistry can do for their appearance. In this case, the authors and the patient chose R-2 L-1 from the Smile Style Guide (Digident, Inc). That means round canines, square round lateral incisors, square centrals, and even length (Figure 11).

### Treatment

All treatment begins with proper records. A caliper was used to measure the ala of the nose (Figure 12) to assist in the placement of the maxillary canines. A papillometer (Figure 13) was used to measure the vestibule to the tip of the lip. The authors took initial impressions of the mandibular denture and the maxillary arch. Because the incisors were so mobile, they were splinted together with light-cured Paint-On Dental Dam (Den-Mat Corporation) before an upper Silgimix

(Sultan Healthcare, Inc) impression was taken. Silgimix, a polyvinyl siloxane alginate substitute, was chosen for its ability to provide excellent flow and detail with minimum tear strength. The authors did not want to remove any teeth with the impression. After the impression, the resin was removed.

A bite registration to relate the opposing occlusal surfaces was made with Genie Bite Fast Set (Sultan Healthcare, Inc). It has been this author's experience that Genie Bite has a consistency that stays where it is placed, does not slump, and approaches almost no resistance to closure. This is very helpful when there are large edentulous areas present, as in the posterior of this case. When set, it does not compress or flex. This helps the dentist position the models with exceptional precision, even without rims.

A maxillary custom tray and bite rim was fabricated for the next appointment. After the liquid rubber dam was placed around the remaining teeth and light-cured, a regular body vinylpolysiloxane (VPS) impression with Genie Impression Material (Sultan Healthcare, Inc) was taken using border molding techniques.

According to the manufacturer, the Genie VPS impression material is chemically engineered to produce remarkably detailed and precise impressions every time. Because of



**Figure 15**—Trimmed Hydro-Cast on immediate.



**Figure 16**—Trimmed Hydro-Cast on immediate, lateral view.



**Figure 17**—Final retracted view.



**Figure 18**—Final smile.

its ultrahydrophilicity, Genie is extremely forgiving in the oral environment. It has been the author's experience that Genie's flow and flexibility gives great detail, and yet the impression can be easily removed from the mouth. This is very important because of the mobility of the teeth involved. Again, the author did not want to remove any teeth unplanned. The author's practice uses this product routinely for removable as well as fixed prosthodontics. The author had also noticed that Genie helps eliminate pulling out restorations along with the impression. The natural occlusion was verified with the bite rim and Genie Bite (Sultan Healthcare, Inc).

### Laboratory Work

In the laboratory, each tooth on the cast, starting from the central incisor and going distal, was cut 2 mm gingival to the cemento-enamel junction and replaced by the artificial one. Teeth were set one at a time. Any slight change in tooth length, even 1 mm to 2 mm, can make the smile line too long or too short. Slight diastemas and overlapping help create a natural effect; however, final decisions should be left to the patient. In this case, he wanted "nice, straight, white teeth." Posterior teeth function better with a slight linguoversion. The immediate transitional denture was then processed for delivery at the surgical appointment.

### Delivery

His maxillary teeth were extracted atraumatically. The immediate transitional denture was relined with Hydro-Cast (Sultan Healthcare, Inc) tissue treatment material (Figure 14). The fingerlike extensions in the sockets were

clipped with surgical scissors and the borders were trimmed with a scalpel (Figures 15 and 16). The immediate denture was delivered (Figures 17 and 18) and the occlusion was adjusted.

It has been this author's experience that tissues in the edentulous mouth react to stresses caused by immediate dentures as well as old, ill-fitting dentures. This results in pain, inflammation, swelling, and tissue displacement (denture stomatitis). This situation is uncomfortable for the patient and can delay the fabrication of new dentures. Hydro-Cast is a temporary soft denture liner that has been proven to reduce the clinical symptoms of denture stomatitis; it helps the tissue return to a state of normal health, and is the only tissue conditioning resin with long-term clinical research demonstrating its efficacy.<sup>8,9</sup> The edentulous tissue surface was recorded in a dynamic functional impression as the gingiva returns to health. Early functional stimulation through the use of a tissue conditioner under an immediate denture is the key to good ridge formation. Tissue covered in this way will generally appear more rounded, smoother, more resilient, and thicker.

Immediate complete dentures contain the edema and mold the tissues for the first 2 days only. The patient was instructed not to disturb the surgical area, rinse vigorously, or probe the area with any objects or fingers. Though this patient was not a smoker, all patients are told not to smoke for at least 72 hours. Smoking can be detrimental to the healing process. It is important to maintain a healthy diet to promote healing, gain strength, have less discomfort, and feel better. For approximately the first 48 hours, he was instructed

to eat a soft diet, such as eggs, yogurt, and soup, to make him feel more comfortable. He was to avoid salty, spicy, or extremely hot foods as well as chips, nuts, popcorn, or anything with sharp edges.

### The Transitional Stage

The patient was made aware that an immediate denture is a trial denture and part of the process. The occlusion was equilibrated to fit the opposing arch. Thanks to the accurate two-bite registrations at the two preliminary appointments, the adjustments were minimal. After an appropriate period of 8 to 12 weeks for healing, the old mandibular denture and new immediate transitional maxillary denture were to be remade. He physically felt better about himself and he loved his new smile. He was even wondering why he needed to return for follow-up treatment.

### Summary

In this article, treatment of a patient with periodontal disease, failing dentition, and an old, ill-fitting mandibular denture were restored with an immediate complete maxillary denture. Part 2 follows next month with the restoration of the oral complex with new complete maxillary and mandibular dentures. ■

### Acknowledgment

The author would like to give special thanks to David Wiley of Denture Technology at 866.253.3038, for his work on this case and so many others.

### References

1. Beck JD, Hunt RJ, Locker D, et al. Variations on the social impact of oral conditions among older adults in South Australia, Ontario, and North Carolina. *J Dent Res*. 1996;75(7):1439-1450.

2. Gregoire ST, Guillory VL, Mitchell MS, et al. The effect of bioadhesion on in-depth disinfection of denture base acrylic resin. *J Calif Dent Assoc*. 1998;26(11):846-850.
3. Chau VB, Saunders TR, Pimsler M. In-depth disinfection of acrylic resins. *J Prosthet Dent*. 1995;74(3):309-313.
4. Cruz GD, Franklin DM, Hay JL, et al. Oral cancer knowledge, risk factors and characteristics of subjects in a large oral cancer screening program. *J Am Dent Assoc*. 2002;133(8):1064-1071.
5. Oral Cancer Foundation. Rates of occurrence in the United States. Available at: [www.oralcancerfoundation.org/facts/index.htm](http://www.oralcancerfoundation.org/facts/index.htm). Accessed Feb. 9, 2006.
6. Oral Cancer Foundation. Alcohol and Tobacco. Available at: [www.oralcancerfoundation.org/facts/alcohol\\_tobacco.htm](http://www.oralcancerfoundation.org/facts/alcohol_tobacco.htm). Accessed Feb. 9, 2006.
7. American Cancer Society. Detailed guide: oral cavity and oropharyngeal cancer. What are the risk factors for oral cavity and oropharyngeal cancer? Available at: [www.cancer.org/docroot/CRI/content/CRI\\_2\\_4\\_2X\\_What\\_are\\_the\\_risk\\_factors\\_for\\_oral\\_cavity\\_and\\_oropharyngeal\\_cancer\\_60.asp](http://www.cancer.org/docroot/CRI/content/CRI_2_4_2X_What_are_the_risk_factors_for_oral_cavity_and_oropharyngeal_cancer_60.asp). Accessed March 12, 2006.
8. De Leeuw R, Boering G, Stegenga B, et al. Clinical signs of TMJ osteoarthritis and internal derangement 30 years after nonsurgical treatment. *J Orofac Pain*. 1994;8(1):18-24.
9. Tassarotti B. A clinical and histologic evaluation of a conditioning material. *J Prosthet Dent*. 1972;28(1):13-18.

### Product References

**Product:** Sea-Bond insert  
**Manufacturer:** Combe Incorporated  
**Location:** White Plains, New York  
**Phone:** 800.873.7400  
**Web site:** [www.combe.com](http://www.combe.com)

**Product:** Smile Style Guide  
**Manufacturer:** Digident, Inc  
**Location:** Dallas, Texas  
**Phone:** 800.741.7966  
**Web site:** [www.digident.com](http://www.digident.com)

**Product:** Paint-On Dental Dam  
**Manufacturer:** Den-Mat Corporation  
**Location:** Santa Maria, California  
**Phone:** 800.445.0345  
**Web site:** [www.denmat.com](http://www.denmat.com)

**Products:** Silgimix, Genie Bite Fast Set, Genie Impression Material, Hydro-Cast, Genie Bite  
**Manufacturer:** Sultan Healthcare, Inc  
**Location:** Englewood, New Jersey  
**Phone:** 800.637.8582  
**Web site:** [www.sultanintl.com](http://www.sultanintl.com)



# Contemporary Esthetics

## Information for Authors:

*Contemporary Esthetics* is seeking articles focused on the latest procedures and technological advances in esthetic and restorative dentistry. If you would like to submit an article for consideration, please refer to the guidelines below. For more information, please contact:

Joseph P. Depa III  
Managing Editor  
241 Forsgate Drive  
Jamesburg, NJ 08831  
800.926.7636 ext. 331  
jdepa@ascendmedia.com

*Contemporary Esthetics* welcomes original, unpublished articles on esthetic and restorative dentistry. Our esthetic Case Studies include a complete patient history, discussion of the treatment planning involved, the step-by-step procedure that was followed, including products used, and a “before and after” evaluation of the patient’s condition.

### ■ Manuscript Format

Each Case Study article should consist of no more than 3,000 words, and should include references, 15 to 25 figures and corresponding captions, and any necessary tables. Make sure all corresponding figures are cited in the text in numerical order. Authors must include figure captions for all figures provided with manuscripts. If an article is accepted as CE, the author will be asked to provide a review quiz with 10 questions, 3 or 4 learning objectives, and a 1-paragraph abstract. Manuscripts may be submitted via email.

A color head shot and author biography (not to exceed 2 paragraphs), including credentials and contact information (city/state, phone, fax, email) for private or other practices, should be included for all authors and coauthors of the manuscript.

Accepted manuscripts will be edited for style, grammar, and format. Authors will have an opportunity to check the progress of their manuscripts or make any additions/corrections before publication.

### ■ Illustrations

Manuscripts should be accompanied by 2 complete sets of 15 to 25 color figures (preferably slides). Write your name, the figure number, and orientation indicators on the back of each photograph, on the front of each slide frame, or on the front of each printout. (Indicate the “front” and “top” of the slide according to how the slide should appear in print, not in a slide projector.)

### ■ Submitting Electronic Files

It is imperative that, if you plan on submitting images electronically, you adhere to the following formats. We accept 3 graphic file formats: jpeg, tiff, and eps. Any images submitted without these extensions will NOT BE ACCEPTED. **DO NOT EMBED IMAGES IN WORD**

**DOCUMENTS. WE CANNOT USE EMBEDDED IMAGES. DO NOT SUBMIT POWERPOINT SLIDE SHOWS.** The only acceptable formats are tiff, jpeg, and eps.

Electronic graphic files must also be high-resolution. This means that the original photo/slide must be scanned at the following settings: 300 dpi (dots per inch); 3 inches in width; 4-color (CMYK). Please adhere to these formats. Images that do not meet these criteria cannot be used and will be returned for rescanning, holding up the editorial process.

Electronic images can be sent via email, or placed on CD or Mac-formatted zip disks. Include a color printout of each image. All original artwork/disks will be returned promptly after manuscript publication.

### ■ References

References/citations must appear in the text as typed, numbered superscripts (not footnote annotations) and should be listed at the end of the manuscript in their order of appearance. References cited should be less than 10 years old. Provide complete bibliographic information for all materials cited. References should be formatted in AMA style.

### ■ Product References

Manuscripts that cite several trade-name products must also include Product References in order to be considered for publication. Each trade-name product mentioned in the article should include the corresponding manufacturer’s name in parentheses.

Additionally, these same products, along with the manufacturer’s name, address, phone, and fax, must also be included in an alphabetical list at the end of the article.

### ■ Manuscript Review

All articles are peer-reviewed by the Editor-in-Chief and members of the editorial board. The review process usually takes 8 weeks. The Editor-in-Chief and publisher reserve the right to make revisions in the text where appropriate. If questions arise during the review process, the author will be contacted.

### ■ Learning Objectives

All CE manuscripts must include specific learning objectives so that readers will be able to identify at the outset what they can expect to learn from the article. Objectives should be phrased using verbs to make clear what the reader will learn, not what the author will discuss. Each objective should be stated in terms that clarify its clinical relevance. Review questions must address stated objectives.

### ■ Review Questions

CE manuscripts must be accompanied by 10 multiple-choice Review Questions (answers must be located directly in the text) so that readers may gain Continuing Education Credits. No true/false questions will be accepted. Questions must have an opening statement, one correct answer, and three distractors. Indicate the correct answer.

### ■ Conflict of Interest Disclosure

*Contemporary Esthetics* requires that all authors disclose their own or a family member’s commercial or financial interest in any related products or developmental or research relationships with companies that manufacture related products.

### ■ Copyright Reproduction Permission

Tables, photographs, or illustrations reproduced exactly from another work must be accompanied by written permission from the original source. Full credit will be given to the original source.

